Journal of Research and Advancement in Dentistry

Volume 12 Number 3 May - June 2021

www.jrad.co.in

Official Publication of Magadh University and Kolhan University

Rehabilitation of Posterior Maxilla Using Implants

Anshuman Chaturvedi, Shitij Srivastava, Abhinav Shekhar, Utkarsh Gupta*

Departement of Prosthodontics, Crown Bridge and Implantology, Sardar Patel Post Graduate Institute of Dental and Medical, Lucknow, Uttar Pradesh, India.

Abstract

Background: Present article describes the various treatment modalities available for posterior maxillary rehabilitation. Preservation of the remains and restoration of the missing is needy to maintain the form and function of the stomatognathic system. Loss of a tooth can be common in any of the arches. The rehabilitation of the lost dental and alveolar part in the maxillary arch is more technique sensitive in comparison to the corresponding arch. The inferior quality of bone in comparison to the mandibular arch fixed restorative procedures is quite cumbersome to the practitioner. This article is an attempt to explain the various options that can be opted for the rehabilitation of the edentulous maxillary arch according to the diagnosed condition of the individual.

Keywords: Implant, Sinus lift, Zygomatic implants, Tuberosity implant, Pterygoid implants.

INTRODUCTION

Today in dentistry, implants play a crucial role.^[1] One of the factors affecting the stability of the implant is bone quality which varies from maxilla to mandible. The posterior maxilla has mostly the D3-D4 quality of bone which affects the primary stability of the bone. The primary stability of the implants in the posterior maxillary region can be attained by the adequate length and width of the implant, which is not possible due to the presence of the maxillary sinus.^[2] Hence, it will direct us to enhance more improvised options and techniques for the rehabilitation of the posterior maxillary region. In the present dissertation, we are discussing the various options to restore the posterior maxilla. As the implant with the normal surgical approach is our priority of time in case of adequate conditions.^[3]

Maxillary sinus

Maxillary sinus (a pyramid-shaped cavity) is the largest of the paranasal air cavities occupying the body of the maxilla and draining into the middle meatus of the ear.^[4] The functional roles of the maxillary or paranasal sinuses continue to be elusive^[5] and include mechanical functions like decreasing the relative weight of the front of the skull especially the bones of the face,^[6,7] providing a buffer against blows to the face and protection to the brain^[7,8] and the function as pillars for the dispersal of masticatory forces,^[9,10] air conditioning, filtering, warming of inspired air for the regulation of intranasal and sinus gas pressures or thermal regulation for the central nervous system,^[11] and phonation by increasing the resonance

of the voice.^[12] The development of the maxillary sinuses begins in the 3rd week of intrauterine life and continues till late adulthood.^[13]

Treatment approach

Misch in 1987 developed a classification for the treatment of edentulous posterior maxilla based on the amount of bone available below the antrum and the ridge width (Figure 1).^[14] Treatment categories ranged from subantral augmentation category 1 (SA1) to SA4 based on height A (>5 mm) and B (2.5–5 mm) based on ridge width.

- SA1: It has an adequate vertical bone for implants, that is, 12 mm. No manipulation of the sinus is required
- SA2: 0–2 mm less than the ideal height of bone in which surgical correction may be required
- SA3: 5–10 mm of bone below the sinus
- SA4: It has less than 5 mm of bone below the sinus.^[14]

Techniques for maxillary sinus augmentation

1. Lateral window approach/direct sinus lift^[15]
2. Transcrestal approach/indirect sinus lift^[16]
3. Minimal invasive surgery

Address for correspondence: Dr. Utkarsh Gupta,

Department of Prosthodontics, Crown Bridge and Implantology, Sardar Patel Post Graduate Institute of Dental and Medical, Lucknow, Uttar Pradesh, India. Email: utkarsh1906@gmail.com

Received: Feb. 13, 2021; **Accepted:** Mar. 02, 2021; **Published:** Apr. 30, 2021

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

How to cite this article: Chaturvedi A, Srivastava S, Shekhar A, Gupta U. Rehabilitation of Posterior Maxilla Using Implants. J Res Adv Dent 2021;12:3:1-5.

Access this article online

Website:
www.jrad.co.in

DOI:
<https://doi.org/10.53064/jrad.2021.12.3.01>

- Piezoelectric surgery
- Hydraulic lift sinus surgery
- Balloon elevation technique
- Magnetic mallet technique

Nasal Suction technique for the lifting of the Schenederian membrane.

Graft versus graft less approach

The criteria for grafting or not grafting the sinus after membrane elevation are the residual bone height and width. If the residual bone height is less than 5 mm, bone grafting may be required, but if it is more than 5 mm, it can be carried out without grafting procedures. If residual bone width is less horizontal, only grafting is done following sinus augmentation, and if it is sufficient, a sinus augmentation can be carried out without bone grafting.

The net bone present is the cumulative effect of bone deposition minus the bone resorption. Coagulum plays a crucial role in endogenous bone formation, and it seems that the osteoinductive properties of a coagulum is limited mainly by lack of a properly maintained space.^[17]

Various treatments options for posterior maxilla

- a. Normal implants (where sinus lift is not required)
- b. Short implants
- c. Zygomatic implants
- d. Tuberosity implants
- e. Pterygoid implants.

NORMAL IMPLANTS

Implant placement in the posterior maxilla is a challenging procedure when vertical deficiencies are present. Sinus membrane elevation plays a crucial role in implant

placement. But sometimes, when the bone height is sufficient (>8mm) then even normal implants can be placed without the need for sinus elevation.^[18]

SHORT IMPLANTS

Given the risk factors, skills needed, and cost of open lateral window sinus grafting, the option to use short or ultra-short dental implants is desirable if the increased risk of implant failure is not an impediment (Table 1). Certainly, press-fit implant designs have shown excellent long-term survival using the transrectal approach. However, many clinicians continue to be intimidated by the use of short endosseous root-form threaded implants — mostly because of widespread peer influence.^[19]

ZYGOMATIC IMPLANTS

The Zygoma Anatomy Guided Approach (ZAGA) developed by Carlos Aparicio is a one-stage procedure for immediate loading. ZAGA classification takes into account the degree of concavity of the anterior maxillary wall and prosthetic implant component. Accordingly, five different anatomical possibilities exist that influence the possible trajectory of zygomatic implants. ZAGA has shown to be better esthetic and prosthetic concern than the Branemark system.

Classification

The five basic anatomical groups were named as (Figure 2)^[20]

1. ZAGA 0
2. ZAGA 1
3. ZAGA 2
4. ZAGA 3
5. ZAGA 4.

Indications for zygomatic implants

1. Extreme resorption in the sinus area
2. Severe atrophy of the posterior maxilla
3. Reconstruction of maxillary defects following tumor resection
4. Unilateral cleft palate and generalized maxillary atrophy
5. Nasal reconstruction.

Advantages of zygomatic implants

1. Avoids use of grafts in atrophic maxilla
2. No additional donor site surgery and morbidity

Figure 1: Subantral treatment categories

Table 1: Advantages and disadvantages of short implants

Advantages	Disadvantages
1. Few surgical procedures are required resulting in less morbidity and less swelling and pain	Achieving initial stability can be challenging.
2. Few materials are required, which means less cost for the patient	Submerged healing with subsequent re-entry is generally recommended
3. Less surgical time	More chances of prosthetic complications
4. Decreased risk of damaging adjacent tooth roots	Type 4 bone is at higher risk

3. Good anchorage from tough zygomatic bone which enhances stability of prosthesis
4. Zygomatic implants do not necessarily require hospitalization, which is usually needed for autogenous bone harvesting from the iliac crest.^[20]

TUBEROSITY IMPLANTS

The maxilla bone tuberosity is the posterior convexity of the maxillary alveolar ridge. The pyramidal process is its medial and posterior boundary. The pyramidal process is located behind and slightly medial to the tuberosity. This process binds to the anterior surface of the pterygoid plates of the sphenoid bone and is interposed between the inferior end of the pterygoid plates and the maxillary tuberosity. The implant is fixed into the pterygoid pillar, which is the junction of palatine bone and pterygoid plates.

If the quality height and volume of maxillary tuberosity are insufficient, a more medially angled and implant placement are determined by the angle of the posterior wall of the maxillary sinus. To prevent penetration of the sinus implant is placed parallel to the posterior sinus wall. Based on the angle of

placement and length, the posterior implant can be classified as follows:

Classification of the posterior maxillary implant based on anatomic location (Figure 3a-d)^[21]

1. Tuberosity
2. Tuberosity/pterygoid process
3. Tuberosity/pyramidal process
4. Tuberosity/pyramidal process/pterygoid process.

If the anatomy of the tuberosity is favorable, a long implant can be placed. Maxillary sinus lining must not be perforated. Tuberosity has a poor bone quality. Hence, a wider implant should be used. The wide-diameter implant fills a greater volume of the tuberosity and provides an osteotome-like effect by condensing and compacting the internal bone of this structure between the implant and the cortical plates of bone.^[21]

PTERYGOID IMPLANTS

Implants placement in the pterygoid region shows proper osseointegration, retention, and stability. It was introduced by Tulasne (1992). Tulasne (1989) credited Paul Tessier for

Figure 2: The five basic anatomical groups

Figure 3: (a) Tuberosity. (b) Implant is located in the tuberosity, and it should be parallel to the posterior wall of the maxillary sinus. (c) Posterior implant engaging the tuberosity and pterygoid process. (d) Posterior implant engaging the tuberosity and pyramidal process

proposing the idea of placing implants in the pterygoid region. The length of pterygoid implants ranges from 15 mm to 20 mm due to a toothier long insertion path. Pterygoid implants have a bicortical anchorage, due to which the axial loading and force dissipation are improved, and the cantilever is eliminated. Pterygoid implants have been defined as “implant placement through the maxillary tuberosity into the pterygoid plate” by the glossary of oral and maxillofacial implants.^[22]

The PARP classification allows working only in the pterygomaxillary region with retromolar implants (Figure 4).^[23]

- PARP 1. It is the simplest scenario when there is no sinus invasion, and we have a bone in all its route. Bone density determines the length of the implant
- PARP 2. The patient presents with a sinus invasion but still has >10 mm of the remaining bone
- PARP 3. This is a case of medium-high difficulty, with sinus invasion leaving a bone surface between 5 mm and 9 mm of remaining bone
- PARP 4. Remaining bone less than 5 mm.

Technique

The pterygoid implant is placed using a tilted concept, called as TTPHIL technique. Implant placement is done in the region of the second and third molar edentulous space toward the junction formed by the posteroinferior projection of the sphenoid, palatine process and the maxillary surface with distal angulation of 25–45 degree depending on the maxillary floor and height of tuberosity.^[23]

COMPLICATIONS ASSOCIATED WITH IMPLANT PLACEMENT IN MAXILLARY POSTERIOR REGION AND ITS MANAGEMENT

Perforations management

In the case of perforation, it is important to evaluate the width and outline of a window to reach an intact membrane from where elevation can be initiated. The membrane surrounding the perforation is delicately dissected with a blunt instrument and the elevation is started far from the perforation in an attempt to relieve the tension on the tissue of the perforated area (Table 2).^[22]

In small perforations (<5 mm), the membrane will fold on itself during the elevation. After complete elevation, an absorbable collagen wound dressing is applied to the antral mucosa as reinforcement. Its adhesive properties make it easy to handle and apply the perforated membrane. A platelet-rich fibrin membrane can also be used as an alternative, followed by bone filling.

In larger perforations (5–10 mm), an absorbable collagen membrane is placed underneath the elevated sinus membrane to create a superior barrier to confine the grafts, followed by suturing the perforation with a 6/0 Vicryl (Ethicon, Norderstedt, Germany) but is a very difficult manipulation because of the inaccessibility and the mechanical properties of the mucosa lining that has a tendency to dilaceration and subsequently to enlarge or create new perforations.^[22]

Complete tear

In this case, a large resorbable collagen membrane allows proper reparation. This membrane will be placed on the medial wall of the sinus and then folded on the borders of the bone window from the upper and lower part as a new roof and floor.

Post-operative care in patients who have undergone sinus grafting, especially with perforation, is very important, and the patient is asked not to blow their noses and not to sneeze, opening the mouth for 2 weeks after surgery (no scuba diving and no flying). If the repair of perforation seems difficult, the procedure is interrupted.^[22]

General considerations

- To achieve primary implant stability (35–45 Ncm insertion torque)
- Indicated with a minimum bone width of 5 mm and minimum bone height of 10 mm from canine to canine in the maxilla and 8 mm in mandible
- If angulation is 30 ° or more, the tilted implants can be splinted
- For tilted posterior implants, the distal screw access holes should be located at the occlusal face of the first molar, the second premolar, or the first premolar.

PARP 1	PARP 2	PARP 3	PARP 4
Without sinus invasion	With or without minimal sinus invasion	Moderate sinus invasion	Critical sinus invasion
Bone >13 mm	10–13 mm remaining bone	5–9.99 mm remaining bone	<5 mm remaining bone
Retromolar/pterygoid	Retromolar/pterygoid	Pterygoid	Pterygoid

PARP: Pterygoid anatomic radiographic prediction			

Figure 4: Classification of the PARP

Table 2: Perforation size, classification, and repair

	Class 1 Perforation	Class 2 Perforation	Class 3 Perforation	Class 4 Perforation	Class 5 Perforation
Size	< 2mm	2 – 5mm	Complete tear	Bony oro-antral communication soft tissue intact	Complete communication, Soft and hard tissue are separated
Repair Technique	Continue with bone grafting	Sinus membrane folding technique	Delayed membrane sandwich technique	Split thickness sinus membrane sandwich technique	Invagination technique

CONCLUSION

Finally, we can conclude that rehabilitation in the posterior maxilla is a very complex procedure due to excessive loss of bone height, nearness to the maxillary sinus, and also lack of availability of space to place implants. Recently several modifications and new techniques are being developed by several companies so as to make sinus elevation much easier. Some of these include the lateral approach sinus kit manufactured by Osstem has a dome drill, a core drill and a stopper to know where to stop the depth of the drill. Another system includes Dentium advanced sinus kit (DASK) system manufactured by Dentium has a special DASK drill and an internal irrigation system so as to cool the heat produced during cutting. There is also another system called crestal approach sinus kit (CAS-KIT) manufactured by Osstem implant system has as specific drill, a hydraulic system to elevate the membrane and special carrier to graft materials. Hiossen's CAS-KIT is specifically designed to easily and safely lift the membrane in the maxillary sinus from a crestal approach. Another system known as PALTOP Advanced Implant DIVA manufactured by Paltop possesses a standard interchangeable hex connection. The PALTOP Standard Platform with a 2.5 mm hexagonal connection covers 3.75, 4.2, and 5.0 mm implant diameters. These three implant diameters utilize one line of prosthetic components.

The ultimate goal in maxillary rehabilitation is to place an implant with less time consumption and less trauma to the patient. Therefore, it is important for a clinician to have a sound knowledge of the maxilla, maxillary sinus, and its various surrounding and associated structures to place a successful dental implant. In all the above, we have tried to discuss and cover all the things starting from anatomy of maxilla and maxillary sinus to various options available to do sinus elevation, various approaches to do direct or indirect sinus lift, various graft materials that can be used and placed and also the various treatment modalities related to posterior maxillary rehabilitation.

CONFLICTS OF INTEREST

The authors declare that they have no potential conflict of interests regarding this article.

REFERENCES

- Hanif A, Qureshi S, Sheikh Z and Rashid H. Complications in implant dentistry. *Eur J Dent* 2017;11:135-40.
- Misch CE, Resnik RR. Misch's Avoiding Complications Oral Implantology. 2nd ed. Maryland Heights, Missouri: Mosby; 2018. p. 16-20.
- Zadeh SR, Ahmad K, Kim H. Prosthetic failure in implant dentistry. *Dent Clin N Am* 2015;59:195-214.
- Asbell MB. Dentistry: A Historical Perspective. Vol. 45. Bryn Mawr, PA: Dorrance-Company; 1988. p. 5660.
- Allen C. Curious Observations on the Teeth. Vol. 687. London. John Bale, Sons and Danielsson, Ltd.; 1687. p. 12-13.
- Ring ME. Pause for a moment in dentistry history: A thousand years of dental implant: A definitive history part 1. *Compend Contin Educ Dent* 1995;16:1060-9.
- Creekmore TD, Eklund MK. The possibility of skeletal anchorage. *J Clin Orthod* 1983;17:266-9.
- Jivraj S, Chee W. Rationale for dental implants. *Br Dent J* 2006;24:661-5.
- Roberts WE, Turley PK, Brezniak N, Fielder PJ. Bone physiology and metabolism. *Calif Dent Assoc J* 1987;15:54-61.
- Devi VS. Inderbir Singh's Human Embryology. 11th ed. New Delhi: Jaypee Brothers Medical Publishers; 2016.
- Sadler TW. Langmans Medical Embryology. 14th ed. Alphen aan den Rijn, Netherlands: Wolters Kluwer; 2021.
- Chaurasia BD. Textbook of Human Anatomy. Head and Neck Part. 6th ed. New Delhi: CBS Publishers; 2017.
- Misch CE. Density of bone: Effect of treatment planning, surgical approach and healing. In: Misch CE, editor. *Contemporary Implant Dentistry*. St Louis: Mosby Publication; 1993. p. 469-85.
- Misch CE. *Contemporary Implant Dentistry*. 3rd ed. St Louis: Mosby; 2007. p. 934-6.
- Tatum H Jr. Maxillary and sinus implant reconstructions. *Dent Clin North Am* 1986;30:207-29.
- Summers RB. A new concept in maxillary implant surgery: The osteotome technique. *Compendium* 1994a;15:152, 154-6, 158.
- Tosta M, Cortes AR, Corrêa L, Pinto DD Jr, Tumenas I, Katchburian E. Histologic and histomorphometric evaluation of a synthetic bone substitute for maxillary sinus grafting in humans. *Clin Oral Implants Res* 2013;24:866-70.
- Thoma DS, Cha JK, Jung UW. Treatment concepts for the posterior maxilla and mandible: Short implants versus long implants in augmented bone. *J Periodontal Implant Sci* 2017;47:2-12.
- Tetradis S, Kantor ML. Extraoral radiographic examination. In: White SC, Pharoah MJ, editors. *Oral Radiology: Principles and Interpretation*. 5th ed. St. Louis, Missouri: Elsevier; 2004. p. 210-24.
- Chow J. *Zygomatic Implants Optimization and Innovation*. Berlin: Springer Publication; 2020.
- Nag PV, Sarika P, Bhagwatkar T, Dhara V. Pterygoid implant: Option for rehabilitation of the atrophic posterior maxilla. *Int J Contemp Dent Med Rev* 2019;2019:011218.
- Proussaefs P, Lozada J. The "Loma Linda pouch": A technique for repairing the perforated sinus membrane. *Int J Periodontics Restorative Dent* 2003;23:593-7.
- Lopes LF, da Silva VF, Santiago JF Jr., Panzarini SR, Pellizzer EP. Placement of dental implants in the maxillary tuberosity: A systematic review. *Int J Oral Maxillofac Surg* 2015;44:229-38.