Periostin a Game-changer in Periodontal Health and Disease: A Review

Priyanka S. Padalkar^{1*}, Manjiri P. Amte¹, Diksha Somkuwar²

¹Department of Periodontology, Aditya Dental College, Beed, Maharashtra, India.

²Department of Prosthodontics, VYWS Dental College and Hospital, Amravati, Maharashtra, India.

Abstract

Background: Periostin is a matricellular protein highly expressed in periosteum, periodontal ligament and is essential for tissue integrity and maturation. It plays a role in collagen fibrillogenesis and is downregulated in periodontal disease. The goal of periodontal regenerative therapy is to predictably restore the tooth's supporting periodontal tissues and form a new connective tissue attachment of periodontal ligament (PDL) fibers and new alveolar bone. Periostin might play a pivotal part in regeneration of the PDL and alveolar bone following periodontal surgery. Recently, several studies have suggested that periostin may be an important regulator of periodontal tissue formation by promoting collagen fibrillogenesis and the migration of fibroblasts and osteoblasts. The aim of this article is to provide an extensive review of the implications of periostin in periodontal tissue biology and its potential use in periodontal tissue regeneration.

Keywords: Periostin, Periodontal ligament, Periodontal tissues.

INTRODUCTION

Chronic periodontitis is an inflammatory disease which is characterized by a progressive gingival inflammatory response to bacterial dental plaque. It leads to clinical attachment loss, pocket formation, alveolar bone loss, increased tooth mobility, and finally the tooth loss.^[1] This condition is characterized by increased secretion of inflammatory mediators such as cytokines and chemokines in the periodontal environment.^[2]

Saliva can also reveal the systemic health as well as the oral health to some extent. Whole saliva may also show the presence of periodontal disease and evidence indicates that the level of some cytokines increases in the saliva of patients with periodontitis. Assessment of the composition of saliva may provide valuable information about biochemical markers for the assessment of periodontal diseases.^[1] Periostin is a hemophilic, secretory protein from the fasciclin I family. It influences cell matrix interactions, cell functions, tissue remodeling, wound repair, and type I collagen fibrillogenesis in periodontal ligament (PDL). It is induced by Transforming growth factor- β (TGF- β) and modulates matrix-cell interactions relevant to connective tissue repair.^[2] Periostin serves as a necessary protein for the integrity, development, and maturity of the tissue and it is believed to play a key role in balancing and regulating homeostasis of the PDL. Expression of periostin is induced

by the tumor growth factor (TGF)- β and bone morphogenetic protein-2 (BMP). It regulates collagen I fibrillogenesis in PDL and reinforces the cross-linking of collagen which improves the mechanical properties of connective tissue of PDL.^[1] Periostin is an extracellular matrix (ECM) protein belonging to the fasciclin family.^[3] It plays a supporting role in cell adhesion, proliferation, survival, and fibrillogenesis in all four components of the periodontium, namely, the PDL, bone, cementum, and gingiva.^[13-17]

HISTORY

Periostin was named in 1999 as a newly discovered factor for matricellular proteins.^[6,18] Periostin is a 90 kDa glutamate-containing secreted matricellular protein. It has 811 amino acids with EMI domain at one end of the chain followed by 4 FAS-1 domains and a carboxyl-terminal domain (Figure 1). There is a C-terminal sequence at the other end of the

Address for correspondence: Priyanka S. Padalkar,

Department of Periodontology, Aditya Dental College, Beed, Maharashtra, India. E-mail: priyapadalkar08@gmail.com

Received: Mar. 8, 2021; **Accepted:** May. 1, 2021; **Published:** Jun. 29, 2021

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online

Website:
www.jrad.co.in

DOI:
<https://doi.org/10.53064/jrad.2021.12.4.13>

How to cite this article: Padalkar PS, Amte MP, Somkuwar D. Periostin A Game-changer in Periodontal Health and Disease: A Review. J Res Adv Dent 2021;12:4:64-68.

Figure 1: Periostin is a member of the fasciclin family

chain.^[15,16] It was first identified in the mouse osteoblastic cell line MC3T3-E1 and initially named osteoblast-specific factor 2 belong to fasciclin-1 family.

Its name is derived from periosteum and PDL, Periostin is preferentially expressed in collagen-rich fibrous connective tissues subjected to constant mechanical stress such as the periosteum, PDLs, tendons, heart valves, and skin. Periostin may be involved in tissue remodeling by promotion of adhesion, cellular differentiation, cell survival, and fibrogenesis.

The periodontium is a specialized tissue that surrounds and also supports teeth, maintaining them in the maxillary and mandibular bones. It provides the support necessary to maintain the function of teeth.

Properties of periostin protein

Periodontium consists of four principal components: Gingiva, PDL, cementum, and alveolar bone. By definition, periodontal tissue engineering/regeneration must achieve regeneration of all tooth-supporting structures, including the alveolar bone, cementum, and PDL, and adequate sealing by the gingival tissue.^[13] Periostin is primarily an inducer of fibrillogenesis and acts by stimulating type I collagen production and cross-linking to generate a tight structure.^[7] It is a fetal protein present in the cervical loop and dental follicle of incisor and molar teeth in cap stage and bell stage, which disappears before tooth eruption. The active role of periostin in tissue regeneration and its potential role in the proliferation of fetal cells related to tooth structure further highlight its significance in morphogenesis and development of bone and periodontal tissue.^[14, 19-22]

Periostin has two main functions namely intracellular fibrillogenesis and extracellular migration.^[23-26] By definition, periodontal tissue engineering/regeneration must achieve the regeneration of all tooth-supporting structures, supporting the reproduction of alveolar bone, cementum, and PDL, and ensuring adequate sealing by gingival tissue.^[13] As periostin is abundantly expressed in the PDL, it may become a potential stock in promoting periodontal tissue regeneration. Its expression is downregulated in human PDL fibroblasts when exposed to tumor necrosis factor alpha and *Porphyromonas gingivalis* lipopolysaccharide both of which are present abundantly in periodontitis.^[7]

Structure of periostin

The periostin protein consists of a signal sequence followed by an EMILIN-like (EMI) domain rich in cysteine, four repeated

and conserved fasciclin-1 (Fas-1) domains, and a C-terminal hydrophilic and variable domain including with other ECM proteins such as collagens, fibronectin (FN), tenascin C, or heparin-binding site. Periostin is classified as a member of the fasciclin 1 family by possessing these typical Fas-1 domains, which involved in cell adhesion functions.^[4, 26-28]

Each Fas-1 domain, rich in glutamate residues, has an N-terminal recognition site for γ -glutamyl carboxylase. It also contains a cell adhesion site that interacts with tenascin-C and BMP-1.^[2,5] In addition, the N-terminal site could recognize the vitamin K-dependent γ -glutamyl carboxylase, to modify glutamic residues to γ -carboxyglutamate post-translationally.^[8] Hence, Gla-containing proteins play important roles in the regulation of coagulation cascade and bone homeostasis.^[29,30]

The EMI domain at the N-terminus is a cysteine residue-rich site for protein-protein interactions through which periostin directly interacts with type-I collagen and FN.^[9,10] The C-terminal domain could alternatively split to generate at least five different human isoforms.^[11]

Functions of periostin

The multiple functions of periostin in dental tissues Figures 4 and 5:

1. Periodontal tissue regeneration: (a) Leading wound healing (b) increase in cell function
2. PDL: (a) collagen structure (b) ECM structure
3. Dental tissue development: (a) Fetal protein
4. Mechanical loading: (a) Mechano-sensory bone and collagen remodeling: The PDL is involved in transmission of forces resulting from mastication. Mechanical strain activates latent TGF- β 1, a known prerequisite for upregulation of certain pro-adhesive and matrix remodeling genes in fibroblasts.^[12,23]

TGF- β 1 is also required for matrix contraction. It, in turn, increases periostin mRNA levels in murine PDL fibroblasts. TGF- β regulates periostin expression not only through activation of the TGF- β receptors but also through the FAK-src signaling pathway, leading to translocation of twist-1 to the nucleus, and subsequent periostin activation.^[12,30]

It has also been seen that removal of mechanical forces or strain inhibited expression of periostin and twist mRNA in the PDL (Afanador *et al.*, 2005). Thus, periostin is essential for maintaining the integrity of PDL fiber system in response to mechanical stress.^[22,28] In a mechanically challenging environment, periostin-null mice showed tissue and cellular defects, and leading to a dramatic loss of periodontal support that, in turn, caused premature tooth loss or pathologic migration.

The cementoblasts also failed to attach to the root surface, thereby resulting in a compromised attachment apparatus. However, in the absence of a mechanically challenging environment (occlusal hypofunction) no periodontal defects developed. Decreasing the mechanical stress corrected the widened PDL and recruited cementoblasts to attach to the root surface.^[25]

POSTN is involved in several cellular and extracellular functions resulting in stimulation of bone formation during development, response to mechanical stress, or bone fracture. Overexpressed in tumors, POSTN may be a potential marker of bone metastasis.

POSTN binds FN through its EMI domain and to BMP-1 through its FAS-1 domain, promoting the deposition of BMP-1 into the matrix. BMP-1 activates LOX precursor (pro-LOX) to mature active form (LOX).^[31-39]

ROLE OF PERIOSTIN IN GINGIVAL CREVICULAR FLUID (GCF), SERUM AND SALIVA

Inflammatory responses to bacteria can initiate the destructive process of periodontitis, leading to both a loss of connective tissue and alveolar bone around teeth and an apical shift of the junctional epithelium. Untreated periodontitis results in loss of function, tissue destruction, loosening, and subsequent loss of teeth.^[40] The preferential ten expression of periostin in collagen-rich tissues submitted to mechanical strains, such as the PDL, and its increase during fracture healing, suggest that it might play a critical role in periodontium maintenance and regeneration. Periostin has successfully been used as a periodontal regeneration marker.^[41,42]

GCF

Padial-Molina *et al.* in ^[43] designed a case-control study to determine the expression profile of periostin that facilitates wound stability and maturation. Periostin increased after periodontal surgery in GCF/wound fluid, which is higher in periodontitis patients. The decline of chronic inflammatory stimuli and bacterial challenge caused by the periodontal surgical procedure could potentially explain the periostin increment. Moreover, the expression levels of periostin in GCF/wound fluid moderated to baseline levels as the wound matured, possibly resulting from an increase in periostin deposits in the ECM as the collagen structure matures.^[43] Kumaresan *et al.*, in 2016, evaluated the level of periostin in GCF of healthy individuals and chronic periodontitis patients who underwent non-surgical periodontal therapy. They showed the level of periostin in the GCF of these can serve as a possible biomarker for the assessment of treatment outcome.^[5]

Saliva

The periostin may be considered as an inflammatory marker in periodontal disease based on the significant relationship between the level of periostin in saliva of chronic periodontitis patients and healthy controls. Eshahrood *et al.* in 2018 conducted the study to compare the concentration of periostin in the saliva of patient with chronic periodontitis and healthy controls in this cross-sectional study. The study showed that periostin was detectable in all samples but its level in saliva of patients with periodontitis was significantly lower than healthy controls.

Periostin in periodontium

The PDL, which lies between the tooth and alveolar bone, is important for many functions such as proprioception, tooth

support, and tissue remodeling in response to physiological or pathological conditions. The PDL consists of cells (*e.g.* fibroblasts, epithelial cells, mesenchymal cells, and bone and cementum cells) and extracellular compartments of dense fibrous connective tissue that confers resistance against mechanical stress. The extracellular compartments are formed of type I, III, and V collagen fibers, FN, and tenascin-C embedded in the intercellular substance Figure 3. It is important for bone and PDL to maintain the local histological architectures and the integrity of the ECM, and several ECM-associated proteins could play pivotal parts in regulating cell proliferation, adhesion, migration, differentiation, survival, *etc.*^[30] Periostin is crucial to maintain the integrity of the PDL. Periostin null mice develop an early onset periodontal disease-like phenotype.^[19]

Periostin is a secreted ECM protein that was originally identified in cells from the mesenchymal lineage (osteoblasts, osteoblast-derived cells, the PDL, and periosteum).^[20,27] It is involved in collagen assembly. Collagen cross-linking is a natural process essential to provide stability to collagen-rich connective tissues. In this process, the two key elements are BMP-1 and lysyl oxidase (LOX). Briefly, BMP-1 cleavages the inactive form of LOX to produce the active LOX enzyme, which, in turn, catalyzes the covalent cross-linking of collagen molecules Figure 2. Interestingly, POSTN binds BMP-1 and collagen I through its FAS1 domains and N-terminal EMI domain, respectively, and thus act as a key player in this process, serving as a scaffold for BMP-1 and collagens to accelerate collagen cross-linking.^[26,32]

The importance of POSTN in collagen cross-linking is also supported by POSTN knockout animal models Figure 6. Thus, POSTN null mice exhibit aberrant collagen fibrillogenesis in the periosteum and a decrease in collagen cross-linking observed in skin, tendons, and heart.^[33]

Figure 2: Properties of periostin protein. Periostin is a 90-kDa secreted protein

Figure 3: Structure of periostin

Figure 4: Multiple function of periostin

Figure 5: Putative functions of periostin in bone

Figure 6: Periostin is a key regulator of collagen cross-linking

Periostin actions have been classified into two major functional categories by Kudo in regard of their molecular properties of protein interactions. First is fibrillogenesis, which occurs inside the cell. And another is cell migration, which including the extracellular action outside the cells.^[36,37] Through its EMI

domain, periostin could interact directly with type I collagen,^[35] FN,^[34] and Notch1,^[37] as well as with tenascin-C and BMP-1^[38] through the Fas-1 domain. These functions fulfill periostin to promote proteolytic activation of lysyl oxidase for collagen crosslinking. Periostin also acts as a ligand of integrins $\alpha\beta 3$ and $\alpha\beta 5$ to facilitate cell motility by activation of actin/myosin contractile machinery.^[39]

CONCLUSION

Periostin holds promise as a reliable inflammatory biomarker for diagnosis and to evaluate the outcome following therapeutic interventions. There is a significant relationship between the level of periostin in saliva and chronic periodontitis. Periostin levels in saliva may be considered as a promising diagnostic potential in periodontal disease.

CONFLICTS OF INTEREST

The authors declare they have no potential conflict of interests regarding this article.

REFERENCES

1. Eshahrood ZR, Vardian ST, Yadegari Z, Admin M, Saravi NS. Periostin levels in saliva of patients with chronic periodontitis. *J Ind Soc Periodontol* 2018;22:25-7.
2. Kumaresan D, Balasundaram A, Naik KV, Appukuttun DP. Gingival crevicular fluid periostin levels in chronic periodontitis patients following nonsurgical periodontal treatment with low level lasertherapy. *Eur J Dent* 2016;10:546-50.
3. Conway SJ, Izuhara K, Kudo Y, Litvin J, Markwald R, Ouyang G, *et al.* The role of periostinin tissue remodeling across health and disease. *Cell Mol Life Sci* 2014;71:1279-88.
4. Kim JE, Kim SJ, Lee BH, Park RW, Kim KS, Kim IS. Identification of motifs for cell adhesion within the repeated domains of transforming growth factor-beta-induced gene, betaig-h3. *J Biol Chem* 2000;275:30907-15.
5. Kii I, Nishiyama T, Li M, Matsumoto K, Saito M, Amizuka N, *et al.* Incorporation of tenascin-C into the extracellularmatrix by periostin underlies an extracellular meshwork architecture. *J Biol Chem* 2010;285:2028-39.
6. Horiuchi K, Amizuka N, Takeshita S, Takamatsu H, Katsuura M, Ozawa H, *et al.* Identification and characterization of a novel protein, periostin with restricted expression to periosteum and periodontal ligament and increased expression by transforming growth factor β . *J Bone Miner Res* 1999;14:1239-49.
7. Kudo A. Periostin in fibrillogenesis for tissue regeneration: periostin actions inside and outside the cell. *Cell Mol Life Sci* 2011;68:3201-7.
8. Coutu DL, Wu JH, Monette A, Rivard GE, Blostein MD, Galipeau J. Periostin, a member of a novel family of Vitamin K-dependent proteins, is expressed by mesenchymal stromal cells. *J Biol Chem* 2008;283:17991-8001.
9. Kii I, Nishiyama T, Kudo A. periostin promotes secretion of fibronectin from the endoplasmic reticulum. *Biochem BiophysRes Commun* 2016;470:888-93.
10. Norris RA, Damon B, Mironov V, Kasyanov V, Ramamurthi A, Moreno-Rodriguez R, *et al.* Periostin regulates collagen fibrillogenesis and the biomechanical properties of connective tissues. *J Cell Biochem* 2007;101:695-711.
11. Hoersch S, Andrade-Navarro MA. Periostin shows increased evolutionary plasticity in its alternatively spliced region. *BMC Evol Biol* 2010;30:1427-32.
12. Takeshita S, Kikuno R, Tezuka K, Amann E. Osteoblast-specific factor 2: Cloning of a putative bone adhesion protein with homology with the

- insect protein fasciclin I. *Biochem J* 1993;294:271-8.
13. Padial-Molina M, Marchesan JT, Taut AD, Jin Q, Giannobile WV, Rios HF. Methods to validate tooth-supporting regenerative therapies. *Methods Mol Biol* 2012;887:135-48.
14. Kruzynska-Freitag A, Wang J, Maeda M, Rogers R, Krug E, Hoffman S, *et al.* Periostin is expressed within the developing teeth at the sites of epithelial-mesenchymal interaction. *Dev Dyn* 2004;229:857-68.
15. Ben QW, Zhao Z, Ge SF, Zhou J, Yuan F, Yuan YZ. Circulating levels of periostin may help identify patients with more aggressive colorectal cancer. *Int J Oncol* 2009;34:821-8.
16. Kudo A. Introductory review: Periostin gene and protein structure. *Cell Mol Life Sci* 2017;74:4259-68.
17. Tang Y, Liu L, Wang P, Chen D, Wu Z, Tang C. Periostin promotes migration and osteogenic differentiation of human periodontal ligament mesenchymal stem cells via the Jun amino-terminal kinases (JNK) pathway under inflammatory conditions. *Cell Prolif* 2017;12:369-72.
18. Du J, Li M. Functions of periostin in dental tissues and its role in periodontal tissues' regeneration. *Cell Mol Life Sci* 2017;74:4279-86.
19. Rios H, Koushik SV, Wang H, Wang J, Zhou HM, Lindsley A, *et al.* Periostin null mice exhibit dwarfism, incisorenamel defects, and an early-onset periodontal disease-like phenotype. *Mol Cell Biol* 2005;25:11131-44.
20. Salvi GE, Lang NP. Diagnostic parameters for monitoring peri-implant conditions. *Int J Oral Maxillofac Implants* 2004;19 Suppl:116-27.
21. Romanos GE, Asnani KP, Hingorani D, Deshmukh VL. Periostin: Role in formation and maintenance of dental tissues. *J Cell Physiol* 2014;229:1-5.
22. Jackson-Boeters L, Wen W, Hamilton DW. Periostin localizes to cells in normal skin, but is associated with the extracellular matrix during wound repair. *J Cell Commun Signal* 2009;3:125-33.
23. Rangiani A, Jing Y, Ren Y, Yadav S, Taylor R, Feng JQ. Critical roles of periostin in the process of orthodontic tooth movement. *Eur J Orthod* 2016;38:373-8.
24. Balli U, Keles ZP, Avci B, Guler S, Cetinkaya BO, Keles GC. Assessment of periostin levels in serum and gingival crevicular fluid of patients with periodontal disease. *J Periodontol Res* 2015;50:707-13.
25. Aral CA, Koseoglu S, Saglam M, Pekbagriyanik T, Savran L. Gingival Crevicular Fluid and Salivary Periostin Levels in Non-Smoker Subjects With Chronic and Aggressive Periodontitis : Periostin Levels in Chronic and Aggressive Periodontitis. *Inflammation* 2016;39:986-93.
26. Coli P, Christiaens V, Sennerby L, Bruyn H. Reliability of periodontal diagnostic tools for monitoring peri-implant health and disease. *Periodontol* 2000 2017;73:203-17.
27. Javed F, Al-Hezaimi K, Salameh Z, Almas K, Romanos GE. Proinflammatory cytokines in the crevicular fluid of patients with peri-implantitis. *Cytokine* 2011;53:8-12.
28. Rios HF, Ma D, Xie Y, Giannobile WV, Bonewald LF, Conway SJ, *et al.* Periostin is essential for the integrity and function of the periodontal ligament during occlusal loading in mice. *J Periodontol* 2008;79:1480-90.
29. Oldenburg J, Marinova M, Muller-Reible C, Watzka M. The Vitamin K cycle. *Vitam Horm* 2008;78:35-62.
30. Merle B, Garnero P. The multiple facets of periostin in bone metabolism. *Osteoporos Int* 2012;23:1199-212.
31. Moussad EE, Brigstock DR. Connective tissue growth factor: What's in a name? *Mol Genet Metab* 2000;71:276-92.
32. Maruhashi T. Interaction between periostin and BMP-1 promotes proteolytic activation of lysyl oxidase. *J Biol Chem* 2010;285:13294-303.
33. Canty EG, Kadler KE. Procollagen trafficking, processing and fibrillogenesis. *J Cell Sci* 2005;118:1341-53.
34. Kaklamanos EG, Tsalikis L. A review on peri-implant crevicular fluid assays potential in monitoring and predicting peri-implant tissue responses. *J Int Acad Periodontol* 2002;4:49-59.
35. Basegmez C, Yalcin S, Yalcin F, Ersanli S, Mijiritsky E. Evaluation of periimplant crevicular fluid prostaglandin E2 and matrix metalloproteinase-8 levels from health to periimplant disease status: A prospective study. *Implant Dent* 2012;21:306-10.
36. Guncu GN, Akman AC, Gunday S, Yamalik N, Berker E. Effect of inflammation on cytokine levels and bone remodelling markers in peri-implant sulcus fluid: a preliminary report. *Cytokine* 2012;59:313-6.
37. Tanabe H, Takayama I, Nishiyama T, Shimazaki M, Kii I, Li M, *et al.* Periostin associates with Notch1 precursor to maintain Notch1 expression under a stress condition in mouse cells. *PLoS One* 2010;5:e12234.
38. Maruhashi T, Kii I, Saito M, Kudo A. Interaction between periostin and BMP-1 promotes proteolytic activation of lysyloxidase. *J Biol Chem* 2010;285:13294-303.
39. Gillan L, Matei D, Fishman DA, Gerbin CS, Karlan BY, Chang DD. Periostin secreted by epithelial ovarian carcinoma is a ligand for alpha(V)beta(3) and alpha(V)beta(5) integrins and promotes cell motility. *Can Res* 2002;62:5358-64.
40. Savage A, Eaton KA, Moles DR, Needleman I. A systematic review of definitions of periodontitis and methods that have been used to identify this disease. *J Clin Periodontol* 2009;36:458-67.
41. Park CH, Rios HF, Jin Q, Sugai JV, Padial-Molina M, Taut AD, *et al.* Tissue engineering bone-ligament complexes using fiber-guiding scaffolds. *Biomaterials* 2012;33:137-45.
42. Panduwawala CP, Zhan X, Dissanayaka WL, Samaranayake LP, Jin L, Zhang C. *In vivo* periodontal tissue regeneration by periodontal ligament stem cells and endothelial cells in three-dimensional cell sheet constructs. *J Periodontol Res* 2017;52:408-18.
43. Padial-Molina M, Volk SL, Rios HF. Preliminary insight into the periostin leverage during periodontal tissue healing. *J Clin Periodontol* 2015;42:764-72.