

## Impaction of Primary Mandibular Canine: A Case Report and Brief Review of the Literature

Manoj Kumar Mallela<sup>1</sup> Surendra Reddy Munnangi<sup>2\*</sup> Uday kumar chowdary Birapu<sup>3</sup> Manasa Poludasu<sup>4</sup> Triveni Titla<sup>5</sup> Aakanksha Vyavahare<sup>6</sup>

<sup>1</sup>Director PG Studies and Professor, Department of Pedodontics and Preventive Dentistry, SVS Institute of Dental Sciences, Mahabubnagar, India.

<sup>2</sup>Reader, Department of Pedodontics and Preventive Dentistry, SVS Institute of Dental Sciences, Mahabubnagar, India.

<sup>3</sup>Senior Lecturer, Department of Pedodontics and Preventive Dentistry, SVS Institute of Dental Sciences, Mahabubnagar, India.

<sup>4</sup>Post Graduate Student, Department of Pedodontics and Preventive Dentistry, SVS Institute of Dental Sciences, Mahabubnagar, India.

<sup>5</sup>Post Graduate Student, Department of Pedodontics and Preventive Dentistry, SVS Institute of Dental Sciences, Mahabubnagar, India.

<sup>6</sup>Post Graduate Student, Department of Pedodontics and Preventive Dentistry, SVS Institute of Dental Sciences, Mahabubnagar, India.

### ABSTRACT

**Background:** Impaction of primary teeth is a rare occurrence and is commonly due to the presence of an odontoma. Odontomas are considered as common odontogenic tumors, which are asymptomatic and found on routine radiographic examination. They appear as radio-opaque lesions of different sizes that are solitary or multiple. Clinically, they are complex or compound, and classified as Intra-osseous where they may erupt but occur inside the bone. Extra-osseous or peripheral occur in the tooth bearing portions of the jaws.

Compound odontomas commonly occur in the anterior region of the upper jaw and complex odontomas in the posterior region of both the jaws. They constitute around 22% of all the odontogenic tumors. Sometimes they might be associated with disturbances in the eruption of teeth like impaction, retention or delayed eruption of primary teeth. This case report presents a rare occurrence of an impacted primary canine with odontoma in a seven-year-old female child.

**Key words:** Odontoma, Primary dentition, Compound odontoma.

### INTRODUCTION

Paul Broca in 1867 coined the term odontoma. Odontomas are developmental anomalies originate from differentiated ectodermal and mesodermal cells of dental lamina from which ameloblasts and odontoblasts arise<sup>1</sup>. In the year 2005, WHO classified them under benign tumors as compound and complex odontomas based on histological and morphological criteria<sup>2</sup>. Compound odontomas is composed of multiple, small tooth like structures and complex odontomas may be amorphous or calcified mass with no morphologic similarity to even a rudimentary teeth. A study of literature reveals occurrence of a few cases of odontoma associated with primary teeth in the mandibular region. They are rarely associated with the primary dentition but commonly present in the

permanent dentition.<sup>3</sup> This case reports an odontoma related to the mandibular region of the primary dentition which is rarely seen.

### CASE REPORT

A seven years old female child reported to the department of Pedodontics and Preventive Dentistry with a chief complaint of asymmetry of face because of a swelling on the right mandibular primary canine region. Family history revealed no signs of unerupted teeth or hypodontia. There was no history of trauma and extra oral examination revealed no submandibular lymphadenopathy on both the sides. The swelling is bony hard in consistency with no crepitations. It is oval in shape extending 2cm antero-posteriorly and 1cm superio-inferiorly.

Received: Feb. 18, 2016; Accepted: Apr. 11, 2016

\*Correspondence Dr. Surendra Reddy Munnangi.

Department of Pedodontics and Preventive Dentistry, SVS Institute of Dental Sciences, Mahabubnagar, India.

Email: surendra.mds@gmail.com

Intra-oral examination revealed the presence of all primary teeth except lower right canine. Past history presents no history of extraction or avulsion of a tooth. There was approximately a space of 4 mm in the mandible between the mesial surface of the first primary molar and distal surface of the primary lateral incisor on the right side. Similar space was absent on the contra-lateral side of the arch. Soft tissue examination showed a non-draining soft tissue swelling with normal overlying mucosa. The swelling was slow growing and reached the present stage (Fig.1a).

A pre-operative radiovisiograph (Fig.1b) and an orthopantomogram (OPG) (Fig.2) revealed an irregularly placed multiple radio-opaque masses similar to the density of a dental tissue.



Fig 1 a&b: Figure showing swelling with preoperative radiograph.



Fig 2: Orthopantomogram.

It had been placed over and around the crown of the impacted primary canine and between the roots of the lateral incisor and the mesial root of the first primary molar. OPG showed a rotated right permanent lateral incisor due to the presence of an impacted primary canine.

A differential diagnosis of an odontoma was made and a surgical profile was advised. The treatment plan was explained to the Parents and an informed consent was obtained prior to the procedure. The treatment was carried out under local anaesthesia by raising a full thickness mucoperiosteal flap. A window was created on the

buccal bone (Fig.3a, 3b) of the primary canine region with a straight fissure bur.




Fig 3 a&b: Window created in buccal bone.

A decision was made to extract the impacted primary canine along with the other 16 denticulate and particulate teeth like structures (Fig.4a), as chances of re-eruption were slim. The cavity was curetted and the sharp bony edges were smoothed with a bony file. The flaps were brought closer and sutured with 3/0 vicryl material (Fig.4b)




Fig 4 a&b: 16 denticulate and particulate teeth like structures removed and sutures placed.

The excised sample was collected and sent for histo-pathological evaluation. Histo-pathological appearance showed presence of tooth like structures resembling pulp tissue in central portion surrounded by dentine and enamel. The report confirmed the diagnosis of a compound odontoma. A space maintainer is planned and the child is under follow up.

## DISCUSSION

"Tooth impaction" refers to the cessation of eruption of a tooth caused by a clinically or radiographically detectable physical barrier in the eruption path or by an ectopic position of the tooth. It is caused by local or systemic factors such as presence of supernumerary teeth, odontomas, cysts, crowded tooth germs, erupted teeth, dilacerations,

ankyloses and systemic genetic inter-relations such as cleidocranial dyostosis and hypopituitarism<sup>4</sup>.

The etiology of odontomas may be due to trauma or infection and inflammatory process, hereditary anomalies (Gardener syndrome, Hermanns syndrome), hyperactive odontoblastic activity and alteration of genetic component, which is responsible for dental development. The present case has not shown any signs of systemic manifestations, hereditary factors or history of trauma. Though WHO has classified under benign odontogenic tumors, they can be regarded as hamartomas or malformations because to its composition and behavior. These malformations once calcified do not develop further and remain in the bone or erupt into the mouth. They are characterized by their slow growth and non-aggressive behavior.

Gravey et al. classified compound odontomas into 3 types. Denticulo type is composed of two or more separated denticles having crown and root, dental hard tissues resembling that of a tooth. Particulate type is composed of two or more separate masses or particles bearing no resemblance to the tooth. The last type is Denticulo-particulate type in which both denticles and particles are present together<sup>5</sup>. Some studies showed more occurrences in males than females, but some authors concluded that there is no gender variation. On the contrary Garcia-consuegra et al. and Hisatomi et al. found a female predilection<sup>6,6</sup>. Literature reveals that the position of the odontoma may occur between the roots in posterior teeth, above the crown, near the apices and between the roots of two teeth<sup>7</sup>.

They are more commonly present in permanent dentition than in primary. The occurrence of them in the primary dentition is more in the maxilla than in the mandible. Compound odontomas are common in the anterior region whereas complex odontomas are more prevalent in the posterior region. Surgery was performed similar to the procedure discussed by Yildirum et al. One of the most common treatments advocated for odontomas in both primary and permanent dentition is surgical removal. If detected at an early stage, they can be removed without disturbing the underlying developing tooth, thereby facilitating

spontaneous eruption, if root formation is incomplete. If the tooth is unerupted, the impacted tooth can be treated with orthodontic traction<sup>8-10</sup>. If insufficient space is present for impacted tooth to erupt, up-righting of the inclined neighboring teeth is necessary.

The chances of re-eruption are more in primary dentition than in permanent dentition. This may be due to superficial placement of impacted tooth, less density of the overlying bone and the presence of succedaneous tooth germ at the apical area, which acts as a physical barrier. Thus, when the obstruction to the eruption pathway is eliminated, the impacted tooth begins to move in the less pressurized gingival direction. Some studies stated that the teeth should be extracted when the chances of spontaneous eruption were slim or when there was no expectation of eruption<sup>11</sup>.

The management depends upon the age of the patient, stage of root development in the impacted primary tooth, status of the developing successor and the adjacent tooth. In the present case, the age of the child was 7 yrs and the root formation was complete. So the chance of spontaneous eruption would be slim. The radiographic findings showed a rotated permanent lateral incisor and an obstructed permanent canine, therefore the primary canine was extracted, and planned for the placement of a space maintainer.

## CONCLUSION

Odontomas cause various problems that include impaction or ectopic eruption if not detected early. They are asymptomatic therefore a routine fully mouth radiographs or orthopantogram will be useful for early diagnosis. The prognosis of the teeth involved with odontoma depends on various factors. Proper treatment planning is necessary to remove or save the tooth involved.

## CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

## REFERENCES

1. Shafer WG, Hine MK, Levy BM. A textbook of oral pathology. 4th ed. Philadelphia: WB. Saunders Company, 1983:308.

2. Kramar IRH, Pindborg JJ, Shear M. World health organization international histological classification of tumours – histological typing of odontogenic tumours. 2nd ed. Berlin Heidelberg: Springer-Verlag, 1992.
3. De Oliveira BH, Campos V, Marcal S. Compound Odontoma diagnosis and treatment: three case reports. *Pediatr Dent*. 2001;23:151-7.
4. Yıldırım G, Tosun G, Kızıloglu D, Durmuş E, Şener Y. An Unusual Association of Odontomas with Primary Teeth. *Eur J Dent*. 2007;1:45-9.
5. Raj K, Shetty SB, Joy A, Shetty RN, Kaikure M. Compound Odontoma: A Case Report. *Int J Adv Health Sci* 2015;1(12):10-13.
6. Salgado H, Mesquita P. Compound odontoma - Case report. *Rev port estomatol med dent cir maxilofac*. 2013;54:3:161-5.
7. Teruhisa U, Murakami J, Hisatomi M, Yanagi Y and Asaumi J. A Case of Unerupted Lower Primary Second Molar Associated with Compound Odontoma. *The Open Dentistry Journal*. 2009;3:173-6.
8. Snawder KD. Delayed eruption of the anterior primary teeth and their management: report of a case. *J Dent Child*. 1974;41:382-4.
9. Otsuka Y, Mitomi T, Tomizawa M, Noda T. A review of clinical features in 13 cases of impacted primary teeth. *Int J Paediatr Dent*. 2001;11:57-63.
10. Kaugars GE, Miller ME, Abbey LM. Odontomas. *Oral Surg Oral Med Oral Pathol*. 1989;67:172-6.
11. Stajcic ZZ. Odontoma associated with a primary tooth. *J Pedod*. 1988;12:415-20.